



Information for you


Published February 2012, Updated October 2022

Corticosteroids in pregnancy to reduce complications from being born prematurely

About this information

This information is for you if you are pregnant and have been offered corticosteroids (commonly known as steroids) because there is a chance that your baby may be born early. It may also be helpful if you are the partner, relative or friend of someone in this situation.

The information here aims to help you better understand your health and your options for treatment and care. Your healthcare team is there to support you in making decisions that are right for you. They can help by discussing your situation with you and answering your questions.



Within this leaflet we may use the terms 'woman' and 'women'. However, it is not only people who identify as women who may want to access this leaflet. Your care should be personalised, inclusive and sensitive to your needs whatever your gender identity.

A glossary of medical terms is available on the RCOG website at:

<https://www.rcog.org.uk/for-the-public/a-z-of-medical-terms/>.

Key Points

- Corticosteroids are a medication that may be offered to you if you are at risk of giving birth early.
- They can increase the chance of your baby surviving if they are born prematurely and can reduce the risk of them having health problems.
- Corticosteroids are of most benefit to your baby if they are given between 24 hours and 1 week of your baby being born.
- Babies born between 24 and 35 weeks benefit most from steroids.

What are corticosteroids?

Corticosteroids are a type of medication that may be offered to you to help your baby if there is a chance that you may give birth early. Steroids are given by an injection into the muscle usually of your thigh or upper arm. A single course can consist of two to four injections usually over a 24–48 hour period.

Why are corticosteroids helpful?

Premature babies may have a range of health problems, which tend to be more serious the earlier they are born.

Giving you steroid injections shortly before your baby is born reduces the risk of them having serious complications including:

- Problems with their breathing
- Bleeding into their brain
- Developmental delay

If steroids are given within seven days of your baby being born prematurely there is a reduced chance of them dying around the time of birth.

Who should be given corticosteroids in pregnancy?

Babies born between 24 and 35 weeks benefit most from steroids.

You will be offered steroids if there is an increased chance that your baby will be born before 35 weeks of pregnancy. This includes:

- if you are in suspected or confirmed premature labour
- if your waters break early even if you are not having contractions (see RCOG patient information on [When your waters break prematurely](#)).
- if it may benefit you or your baby for them to be born early. This may be because your baby is not growing well or because you have developed a problem in your pregnancy such as severe pre-eclampsia (see RCOG patient information on [Pre-eclampsia](#)).

If you are likely to give birth between 35 weeks and 37 weeks, or are having a planned caesarean birth before 39 weeks you should have an individualised discussion with your healthcare professional about the risks and benefits of steroids for your baby. There is less evidence that steroids are helpful after 35 weeks.

The benefits of steroids in twin and triplet pregnancies are less certain, but it is still recommended that should be offered steroids if you are likely to give birth early.

Steroids may be given if your baby is expected to be born between 22 and 24 weeks, but this will require an individualised discussion with your healthcare team about the risks to your baby of being born so prematurely.

How long are corticosteroids effective for?

Steroids are of most benefit if the last dose is given to you between 24 hours and 1 week before the birth of your baby. There may still be some benefit even if your baby is born within 24 hours of the first dose.

The benefits of steroids are likely to be significantly reduced if your baby is born more than 7 days after the treatment. Therefore, it is important to try to give steroids at the right time.

Can corticosteroids harm me or my baby?

A single course of steroids is safe for you. You may experience some minor side effects such as pain at the injection site. If you have pre-existing or gestational [diabetes](#) steroids can affect your blood sugar control. You can find further information on RCOG patient information on [Gestational Diabetes](#).

If you have diabetes, you will need increased monitoring of your blood sugar levels while you have the steroids and may need to be admitted to hospital to be offered additional [insulin](#) treatment.

Treatment with steroids between 22 and 35 weeks pregnant is likely to be safe and beneficial for your baby. No long-term harm has been shown although there have been no large studies.

Some research suggests that there may be an impact on mental wellbeing later in life for babies born at full term (after 37 weeks) who were given steroids between 22 and 37 weeks.

If you are given steroids and are more than 35 weeks pregnant there is a chance that your baby may have low blood sugar levels after they are born. Low blood sugar can be harmful for babies if it is not treated and can mean your baby needs to be admitted to the [neonatal unit](#). Your baby will be monitored for this and offered treatment if needed.

Small studies suggest that if you are given steroids before a planned caesarean birth after 37 weeks of pregnancy there is a possible link between this and your child not doing as well at school when they are older.

Can I have more than one course of corticosteroids in this pregnancy?

If you have had a course of steroids and you do not give birth in the next 7 days then a second course might have some beneficial effects on your baby's breathing if they are still expected to be born prematurely. Evidence to support this benefit is limited and there is also evidence that a repeat course of steroids may mean your baby is smaller than they should be when they are born. Your healthcare professional will advise you of the risks and benefits of a repeat course of steroids depending on your individual situation.

When are corticosteroids not necessary?

If you are unlikely to give birth in 7 days, you should not receive steroids. Even if you are at higher risk of giving birth early you

won't be given steroids unless you are likely to give birth in the next 7 days.

Further information

RCOG patient information, When your waters break early:

<https://www.rcog.org.uk/for-the-public/browse-all-patient-information-leaflets/when-your-waters-break-prematurely/>

RCOG patient information, Having more than one baby (Multiple pregnancy):

<https://www.rcog.org.uk/for-the-public/browse-all-patient-information-leaflets/multiple-pregnancy-having-more-than-one-baby/>

RCOG patient information, Gestational diabetes:

<https://www.rcog.org.uk/for-the-public/browse-all-patient-information-leaflets/gestational-diabetes/>

RCOG patient information, Pre-eclampsia:

<https://www.rcog.org.uk/for-the-public/browse-all-patient-information-leaflets/pre-eclampsia-patient-information-leaflet/>

NICE guideline on Preterm labour and birth:

www.nice.org.uk/guidance/ng25

Little Heartbeats: <https://www.little-heartbeats.org.uk/>

Best Beginnings: <https://www.bestbeginnings.org.uk/>

Bliss: <https://www.bliss.org.uk/>

Making a choice

Making a choice

Ask 3 Questions

If you are asked to make a choice, you may have lots of questions that you want to ask. You may also want to talk over your options with your family or friends. It can help to write a list of the questions you want answered and take it to your appointment.



1. What are my options?
2. How do I get support to help me make a decision that is right for me?
3. What are the pros and cons of each option for me?

*Ask 3 Questions is based on Shepherd HL, et al. Three questions that patients can ask to improve the quality of information physicians give about treatment options: A cross-over trial. Patient Education and Counselling, 2011;84: 379-85

<http://aqua.nhs.uk/resources/shared-decision-making-case-studies/>

Sources and acknowledgements

This information has been developed by the RCOG Patient Information Committee. It is based on the RCOG guideline ***Antenatal corticosteroids to reduce neonatal morbidity and mortality*** (Published February 2022). It contains a full list of the sources of evidence we have used. You can find it online at: <https://www.rcog.org.uk/guidance/browse-all-guidance/green-top-guidelines/antenatal-corticosteroids-to-reduce-neonatal-morbidity-and-mortality-green-top-guideline-no-74/>.