



Royal College of
Obstetricians &
Gynaecologists

Guidance for rationalising early pregnancy services in the evolving coronavirus (COVID-19) pandemic

Information for healthcare professionals

Version 1.2: Published Friday 15 May 2020



Summary of updates

Version	Date	Summary of changes
1.1	21.04.20	5.1: Added 'Evidence of a septic miscarriage - signs of infection (e.g. temperature, offensive smelling discharge) in association with symptoms of retained pregnancy tissue (pain and/or bleeding)' as a reason for assessment within 24 hours. Added additional risk factors for ectopic pregnancy.
1.1	21.4.20	10: Section and recommendation added 'Administer anti-D prophylaxis to women who have a surgical procedure, including manual vacuum aspiration, or have a late miscarriage, in line with British Society of Haematology and NICE guidelines.'
1.2	15.05.20	1: Section on BAME advice added 'When reorganising services, units should be particularly cognisant of emerging evidence that black, Asian and minority ethnic group (BAME) individuals are at particular risk of developing severe and life-threatening COVID-19. Clinicians should encourage women to seek early advice if they are concerned about symptoms suggestive of COVID-19. There is extensive evidence on the inequality of experience and outcomes for BAME women during pregnancy in the UK. Particular consideration should be given to the experience of women of BAME background and of lower socioeconomic status, when evaluating the potential or actual impact of any service change.'

I. Introduction

This guidance is to support early pregnancy services during the evolving COVID-19 pandemic. It outlines which elements of care should be prioritised and recommends modifications to early pregnancy care, given national recommendations for social distancing of pregnant women.

When reorganising services, units should be particularly cognisant of emerging evidence that black, Asian and minority ethnic group (BAME) individuals are at particular risk of developing severe and life-threatening COVID-19. This evidence is detailed in the [RCOG coronavirus in pregnancy guidance](#). Clinicians should encourage women to seek early advice if they are concerned about symptoms suggestive of COVID-19. There

is extensive evidence on the inequality of experience and outcomes for BAME women during pregnancy in the UK. Particular consideration should be given to the experience of women of BAME background and of lower socioeconomic status, when evaluating the potential or actual impact of any service change.

2. Screening of women presenting to early pregnancy services

All women should be asked to attend appointments alone or as per local visiting restrictions during the COVID-19 pandemic.

Where a woman requires a consultation due to the need for physical examination or a scan, a system should be in place for evaluating whether she has [symptoms that are suggestive of COVID-19, or if she meets the current 'stay at home' guidance](#). For similar advice in Scotland, [see here](#). This may be a telephone call before the appointment or an assessment at entry to the department.

If a woman attends an appointment but describes symptoms, she should be advised to return home immediately if clinically stable. A member of clinical staff should then make contact with the woman to risk assess whether an urgent modified appointment is required, or whether the appointment can be conducted via telephone consultation.

If an urgent assessment in person or ultrasound scan is required for a woman with confirmed or suspected COVID-19 infection, a room and an ultrasound machine should be designated for this.

All women with a possible COVID-19 infection must be highlighted to all members of the gynecology, maternity, nursing and anaesthetic teams. If the woman requires admission to hospital, the location will depend on the reason for admission and local policy, until COVID-19 testing confirms her status.

3. Delaying appointments where appropriate

3.1 Pre-existing appointments

A review of the clinical urgency of currently held appointments should be made by the clinical team and women will be contacted as necessary.

3.2 In home isolation for suspected or confirmed COVID-19

If delay is clinically appropriate (Table 5.1), care should be provided via a telephone consultation. If urgent care is required, attendance to hospital should be preceded by a phone call to alert the local unit.

3.3 Rebooking appointments

The local service should decide how best to manage rebooking of appointments (blood tests and/or scans) and the woman should be informed of her new appointment.

3.4 Failsafe

A local failsafe should be established to ensure that appointments for all women are reviewed and, if reoffered, that they are attended. Follow local protocols for follow up of women who do not attend.

4. Coordinating your local early pregnancy unit

As well the usual day-to-day requirements for running an early pregnancy unit, we recommend the following:

- Managers should be aware that staff (or members of their family) may become unwell during the pandemic; daily review of the case load, staffing and contingency planning is advised.
- If a pregnant woman is diagnosed with COVID-19, this should be reported to the [UK Obstetric Surveillance System](#).

- Multi-disciplinary team (MDT) meetings: we highly encourage units to conduct a minimum of a weekly MDT meeting (can be arranged using an online meeting platform).

5. Ensuring that early pregnancy units are used appropriately

Women should not attend early pregnancy units without a telephone triage consultation with an experienced clinician, using a locally agreed structure for triage.

Local units following a walk-in model should adopt a robust triage-based system with a dedicated phone number for referrals. Appropriate triage is essential to allow prioritisation of those at high risk of complications, mainly ectopic pregnancy, where hospital visits will be safer than telephone-based consultations.

The inevitable reduction in resources and capacity, as well as the aim to minimise hospital attendance for social distancing of pregnant women, have led to a recommendation of one of the following three options (Table 5.1):

- Scans and/or visits that need to be undertaken without delay;
- Scans and/or visits that can be delayed without affecting clinical care;
- Scans and/or visits that can be avoided for the duration of the pandemic.

Table 5.1 Recommended triage and action for early pregnancy units

Problem	Recommended action
Abdominal or pelvic pain (no previous scan)	Offer scan within 24 hours
Heavy bleeding for more than 24 hours and systemic symptoms of blood loss	Offer scan within 24 hours
Evidence of a septic miscarriage - signs of infection (e.g. temperature, offensive smelling discharge) in association with symptoms of retained pregnancy tissue (pain and/or bleeding).	Offer assessment within 24 hours (Note a temperature may also be associated with COVID-19 infection.)

<p>Pain and/or bleeding together with pre-existing risk factors for ectopic pregnancy:</p> <ul style="list-style-type: none"> • Previous ectopic pregnancy • Previous fallopian tube, pelvic or abdominal surgery, • History of sexually transmitted infections / pelvic inflammatory disease • Use of an IUCD or IUS • Use of assisted reproductive technology • Current smoker or age over 40 	<p>Offer scan within 24 hours if location of pregnancy not known</p>
<p>Moderate bleeding</p>	<p>Telephone consultation with experienced clinician – urine pregnancy test (UPT) in one week:</p> <ul style="list-style-type: none"> • Negative – no follow-up • Positive – offer telephone consultation +/- repeat UPT in one further week or scan
<p>Heavy bleeding that has resolved</p>	<p>Telephone consultation with experienced clinician – UPT in one week:</p> <ul style="list-style-type: none"> • Negative – no follow-up • Positive – offer telephone consultation +/- repeat UPT in one further week or scan
<p>Reassurance</p>	<p>Telephone consultation with experienced clinician – no routine scan</p>
<p>Previous miscarriage(s)</p>	<p>Telephone consultation with experienced clinician – no routine scan</p>
<p>Light bleeding with/without pain that is not troublesome to patient</p>	<p>Telephone consultation with experienced clinician – no routine scan</p>

6. Management of miscarriage

Women who experience a miscarriage should be cared for in accordance with local protocols. There should be an effort to reduce inpatient admission due to COVID-19: offer expectant management for incomplete miscarriage and consider medical management / use of manual vacuum aspiration for missed miscarriage.¹ Counselling should be offered and performed over the phone where possible.

The availability of surgery will need to be reviewed locally on a daily basis and if surgical management is indicated, appropriate precautions related to personal protective equipment (PPE) should be taken in line with national [Health Protection guidance](#).²

Regional anaesthesia may be considered in COVID-19 positive women to reduce the risk to staff from general anaesthetic, which is an aerosol-generating procedure.

Outpatient management is preferred where appropriate. Provide advice on analgesia and the process of miscarrying, in order to support women to remain at home.

Those who have expectant or medical management should not be offered further routine ultrasound scans but asked to repeat a hCG urine test after three weeks. If this is positive, they should be advised to call the early pregnancy unit to arrange further care.

Units should aim to provide telephone consultation to women three weeks following their miscarriage to assess physical and emotional well-being, if resources are available.

7. Intrauterine pregnancy of unknown viability

No further ultrasound scans are recommended.

If the ultrasound scan findings are consistent with menstrual dates no follow up is required.

If findings are not consistent with menstrual dates, explain the risk of miscarriage and consider telephone follow-up in two weeks.

8. Management of pregnancy of unknown location

Use serial beta human chorionic gonadotrophin (beta-hCG) monitoring +/- progesterone at presentation, as per local protocol, to triage women into one of:

- Low risk failing PUL:
 - Pregnancy test at home in two weeks
 - Contact local unit if positive
- Low risk intrauterine pregnancy:
 - Scan in one week to confirm location and viability
- High risk for ectopic pregnancy:
 - return for a repeat beta-hCG and/or scan in a further 48 hours

The M6 model can be used to help with decision making in women with PUL to reduce the number of hospital visits due to COVID-19. It is available at www.earlypregnancy care.co.uk.^{3,4}

9. Management of ectopic pregnancy

Women with ectopic pregnancy should be cared for in accordance to local protocols with an emphasis on conservative management if possible.

9.1 Expectant management

Ensure follow up is appropriate with an individualised approach. There is a need to balance safety with reducing hospital attendance as much as possible in order to reduce the risk of COVID-19 to women, staff and other patients.

When performing beta-HCG monitoring, where possible, repeat levels on a weekly basis. Repeat ultrasound scans should not be routine unless clinically indicated.

9.2 Medical management with single dose methotrexate

It is likely the detrimental effects of methotrexate in COVID-19 are minimal in well women.

As with any ectopic pregnancy, women with suspected /confirmed COVID-19 should be discussed at the early pregnancy unit multi-disciplinary team (MDT) meeting. Administration of methotrexate must be discussed and signed off by a senior clinician prior to treatment, and any ultrasound and beta-hCG levels reviewed carefully. Severely unwell women with COVID-19 and ectopic pregnancy will need to be discussed at an MDT with medical and anaesthetic input.

In addition to routine information giving when offering the choice of methotrexate, inform the woman that:

- Methotrexate is a mildly immunosuppressive medication but there is not thought to be a significant risk in the case of COVID-19 at the dose used to manage ectopic pregnancy. ¹
- There is a theoretical risk that any immunosuppressive medication can make you more vulnerable to viral illness.
- Expert opinion is that the dose of methotrexate given for medical management of ectopic pregnancy is unlikely to increase vulnerability to COVID-19 and does not require home shielding after administration.
- Medical management of ectopic pregnancy avoids hospital admission and surgery, potentially lowering overall exposure to COVID-19.

9.3 Surgical management

Surgical management of ectopic during the coronavirus pandemic should only be considered following senior review of the ultrasound scan, beta-hCG and clinical findings and if no other management option is safely feasible.

The BSGE/RCOG support the use of laparoscopy, but with necessary caution.⁵ Given the limited evidence on the safety of laparoscopy, any laparoscopic surgery should only be undertaken with strict precautions taken

to filter any CO₂ escaping into the operating theatre and the theatre staff wearing appropriate PPE. Mini-laparotomy can be considered as an alternative to laparoscopy if these strict precautions cannot be confidently met.

10. Anti-D prophylaxis

Administer anti-D prophylaxis to women who have a surgical procedure, including manual vacuum aspiration, or have a late miscarriage, in line with British Society of Haematology⁶ and NICE⁷ guidelines.

If miscarriage occurs at home, and having to check RhD status would require an additional visit for the woman, it could be omitted if the risk from COVID-19 outweighs the benefit of receiving anti-D immunoglobulin. Providers should discuss the absence of evidence with women and engage in shared decision making.

11. Management of nausea and vomiting in pregnancy

If a woman has nausea and vomiting in pregnancy, she should be assessed over the phone using the PUQE scoring system and advised regarding anti-emetics, as per local protocol.⁸ Local arrangements for issuing prescriptions remotely after a telephone consultation, where these do not already exist, should be put in place.


Services should plan how to best configure their local protocols during the coronavirus pandemic for those women who require parenteral hydration. This might include hospital at home, day-case or inpatient admission services. Vomiting is a potential risk for transmission, and appropriate.

The rare possibility of a molar pregnancy should be considered in women with hyperemesis gravidarum and other symptoms such as vaginal bleeding. In the event of routine dating ultrasound assessments being delayed, women should be offered assessment in early pregnancy departments if gestational trophoblastic disease is suspected.

References


1. National Institute for Health and Care Excellence. Ectopic pregnancy and miscarriage: diagnosis and initial management. Guideline. NICE. 2019
2. COVID-19: Infection, prevention and control guidance 2020 Available from: <https://www.gov.uk/government/publications/wuhan-novel-coronavirus-infection-prevention-and-control/wuhan-novelcoronavirus-wn-cov-infection-prevention-and-control-guidance> [Accessed 05 January 2020].
3. Bobdiwala S, Saso S, Verbakel JY, Al-Memar M, Van Calster B, Timmerman D, Bourne T. Diagnostic protocols for the management of pregnancy of unknown location: A systematic review and meta-analysis. *BJOG*. 2019 Jan;126(2):190-198, doi: 10.1111/1471-0528.15442.
4. Bobdiwala S, Christodoulou E, Farren J, Mitchell-Jones N, Kyriacou C, Al-Memar M, Ayim F, Chohan B, Kirk E, Abughazza O, Guruwadahyhalli, Guha S, Vathanan V, Bottomley C, Gould D, Stalder C, Timmerman D, Van Calster B, Bourne T. Triaging women with pregnancy of unknown location using two-step protocol including M6 model: clinical implementation study. *UOG*. 2019 August 6, doi: 10.1002/uog.20420.
5. British Society of Gynaecological Endoscopy. Joint RCOG/BSGE statement on gynaecological laparoscopic procedures and COVID-19. 2020
6. H. Qureshi, E. Massey, D. Kirwan, T. Davies, S. Robson, J. White, J. Jones, S. Allard. BCSH guideline for the use of anti-D immunoglobulin for the prevention of haemolytic disease of the fetus and newborn. 2014 doi: 10.1111/tme.12091
7. National Institute for Care Excellence. NICE Guideline 126: Ectopic pregnancy and miscarriage: diagnosis and initial management. Apr 2019. Available from: <https://www.nice.org.uk/guidance/ng126/chapter/Recommendations#anti-d-rhesus-prophylaxis> [Accessed 13 Apr 2020]
8. Royal College of Obstetricians and Gynaecologists. The Management of Nausea and Vomiting of Pregnancy and Hyperemesis Gravidarum Green top guideline No 69. 2016 Available from <https://www.rcog.org.uk/globalassets/documents/guidelines/green-top-guidelines/gtg69-hyperemesis.pdf>.

Appendix I: Summary



Appendix 2: Guidance for management of early pregnancy complications during COVID-19 pandemic

PUV - Pregnancy of unknown viability
 PUL - Pregnancy of unknown location
 EPU - Early Pregnancy Unit
 PPE - Personal protective equipment
 UPT - Urinary pregnancy test
 MVA - Manual vacuum aspiration



Authors

Tom Bourne, Imperial College, AEPU, Tommy's National Centre for Miscarriage Research and ISUOG

Chris Kyriacou, Imperial College, Tommy's National Centre for Miscarriage Research

Arri Coomarasamy, University of Birmingham and Tommy's National Centre of Miscarriage Research

Emma Kirk, Royal Free Hospital, AEPU and ESHRE early pregnancy SIG

George Condous, University of Sydney and ASUM

Mathew Leonardi, University of Sydney

Maya Al-Memar, Imperial College and Tommy's National Centre of Miscarriage Research

Rachel Small, Birmingham Heartland Hospital and AEPU

Eddie Morris, RCOG

Pat O'Brien, RCOG

Gemma Goodyear, RCOG Obstetric Fellow

Jen Jardine, RCOG Obstetric Fellow

Sophie Relph, RCOG Obstetric Fellow

DISCLAIMER: The Royal College of Obstetricians and Gynaecologists (RCOG) has produced this guidance as an aid to good clinical practice and clinical decision-making. This guidance is based on the best evidence available at the time of writing, and the guidance will be kept under regular review as new evidence emerges. This guidance is not intended to replace clinical diagnostics, procedures or treatment plans made by a clinician or other healthcare professional and RCOG accepts no liability for the use of its guidance in a clinical setting. Please be aware that the evidence base for COVID-19 and its impact on pregnancy and related healthcare services is developing rapidly and the latest data or best practice may not yet be incorporated into the current version of this document. RCOG recommends that any departures from local clinical protocols or guidelines should be fully documented in the patient's case notes at the time the relevant decision is taken.

 @RCObsGyn  @rcobsgyn  @RCObsGyn



Royal College of
Obstetricians &
Gynaecologists

Royal College of Obstetricians and Gynaecologists, 10-18 Union Street, London, SE1 1SZ

T: +44 (0) 20 7772 6200

E: covid-19@rcog.org.uk

W: rcog.org.uk

Registered Charity No. 213280

My doctor has confirmed that my appointment will take place at the hospital. Can I take someone with me?

Your healthcare team will be able to advise you on your individual situation. You may be asked to come on your own, or with just one accompanying person. Unfortunately, hospitals are unable to allow children to attend with you during this time.

Will I be able to receive an ultrasound scan?

Unfortunately, early pregnancy units will not be able to offer ultrasound scans to provide reassurance to women who have no symptoms. This will also be the case even if you have a previous history of miscarriage or ectopic pregnancy.

For further information and advice in relation to ectopic pregnancy, please see the further information section at the foot of this document.

What will happen if I do experience a miscarriage?

In the unfortunate event of a miscarriage, your care will depend on your individual situation.

- In most cases, it is likely that you will be asked to miscarry naturally without intervention. Your healthcare team will make arrangements with you to ensure you can contact them should you have any concerns during your recovery.
- You may alternatively receive outpatient medical management. Your healthcare team will make arrangements with you to ensure your medical management is monitored.
- Where it is necessary, you may be asked to attend for surgery. Your healthcare team will discuss this with you directly.

For further support and information in relation to miscarriage, please see the further information section at the foot of this document.

What if I am experiencing pregnancy related nausea and vomiting?

If you are experiencing nausea and vomiting in early pregnancy, you should inform your healthcare team, so that they can arrange the right care for you.

Your care will depend on the level and impact of your symptoms. You may be offered anti-emetics (anti-sickness medicine) as well as outpatient treatment including intravenous fluids. It is unlikely you will be admitted to hospital in this situation, unless your symptoms become serious. See further information section at the foot of this information.

Key points for you

- It is important to know that if you experience any health issues during early pregnancy that require you to be seen by your healthcare professional, an appointment in an early pregnancy unit will be offered and you will receive the care you need.
- Whilst hospitals are trying to minimise people entering in order to reduce the spread of the COVID-19 virus and to limit the impact on services, they are organised in such a way that they are able to provide all acute services.
- If you have symptoms that may be associated with miscarriage or ectopic pregnancy, it is very important that you contact your healthcare professional. You will be able to speak with an experienced member of your healthcare team on the phone before your appointment. They will be best placed to advise you as to whether a visit to the hospital is necessary and to ensure you receive the care that you need.

Further information for you

You can find further information on the matters mentioned in this information at the following organisation websites:

- [Ectopic Pregnancy Trust](#)
- [Miscarriage Association](#)
- [RCOG Information for you: Hyperemesis Gravidaram \(Pregnancy Sickness, nausea and vomiting\)](#)
- [Pregnancy Sickness Support](#)

You can also find all the latest guidance and information on how to protect you and your loved ones during COVID-19 at the following organisation websites:

- [Joint guidance](#) from the Royal College of Obstetricians and Gynaecologists, the Royal College of Midwives, Royal College of Paediatrics and Child Health, Public Health England and Health Protection Scotland
- [UK Government guidance explaining social distancing and self-isolation](#)
- [NHS 111 website](#)
- [NHS Inform in Scotland](#)
- [Public Health England](#)
- [Health Protection Scotland](#)
- Mental Health Support